



Letter to the Editor

Non-*Helicobacter Pylori* *Helicobacter* Species in Humans: What Do We Know and Where Should We Go?

Romina Rezaei¹, Alireza Sadeghi¹, Maryam Hasanzarrini², Hossain Salehi³, Amir Mohammad Salehi⁴¹Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran²Clinical Research Development Unit of Shahid Beheshti Hospital, Hamadan University of Medical Science, Hamadan, Iran³Farhikhtegan Medical Convergence Sciences Research Center, Farhikhtegan Hospital, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran⁴Fetal & Pediatric Cardiovascular Research Center, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

Cite this article as: Rezaei R, Sadeghi A, Hasanzarrini M, Salehi H, Salehi AM. Non-*Helicobacter pylori* *Helicobacter* species in humans: what do we know and where should we go? Middle East J Dig Dis 2026;18(2):112-113. doi:10.34172/mejdd.3375

Received: November 25, 2025, **Revised:** February 11, 2026, **Accepted:** February 12, 2026, **ePublished:** April 30, 2026

Dear Editor,

Shortly after the discovery of *Helicobacter pylori* (*H. Pylori*) by Marshall and Warren in 1982, other *Helicobacter* species have been isolated from the human stomach. These gram-negative spiral-shaped non-*Helicobacter pylori* *Helicobacter* (NHPH) species mostly colonize animals, including pigs, dogs, and cats. However, five NHPH species have also been identified in the human stomach, including *H. suis*, *H. felis*, *H. bizzozeronii*, *H. salomonis*, and *H. heilmannii* s.s. Among these, *H. suis*, which naturally colonizes pigs and non-human primates, is the most frequently detected NHPH in humans. NHPHs transmission from an infected animal host to a human occurs through direct or indirect contact with the animal reservoir, such as dogs, pigs, and cats. Additionally, *H. suis* could be transmitted via the consumption of undercooked or raw pork meat.^{1,2}

While *H. pylori* is a well-established gastric pathogen, affecting more than half of the world's population and responsible for a wide spectrum of gastric disorders, NHPHs are detected in a significant proportion of patients with chronic gastritis, peptic ulcers, and mucosa-associated lymphoid tissue (MALT) lymphoma, especially in cases of negative *H. pylori* infection. Studies report varying prevalence rates of gastric NHPHs among patients. While studies report a 0.2%-6% prevalence of gastric NHPHs in unselected symptomatic populations, more recent research from Western countries and Japan indicates a significantly higher prevalence—exceeding 20%—specifically among patients who are *H. pylori*-negative.^{1,3,4}

Due to the overlapping clinical symptoms (epigastric pain, dyspepsia, nausea, vomiting) and histological findings, clinical differentiation between NHPH and *H.*

pylori is challenging.⁵ Endoscopic findings of NHPH species are not as widely known as those of *H. pylori*. However, advances in image-enhanced endoscopy (IEE), such as linked-color imaging (LCI) and blue-laser imaging (BLI), have improved the visibility and detection of characteristic features. Several endoscopic features, such as white marbled appearance, cracked-like mucosa, spotty redness, and nodular gastritis-like appearance, have been frequently reported in cases of NHPH gastritis.⁶

Traditional histological methods for detecting *H. pylori* have limited diagnostic accuracy in detecting and differentiating NHPHs due to morphological resemblance and low bacterial load. Therefore, targeted molecular diagnostic methods, such as polymerase chain reaction (PCR), are used to detect *Helicobacter* spp. DNA and differentiate among species with high sensitivity and specificity.⁵ Other diagnostic methods currently used to identify NHPHs include histopathological and immunohistochemical findings, culture tests, and enzyme-linked immunosorbent assays.⁷

At present, there is no standardized guideline for eradicating NHPH, and treatment approaches are often empirically based on regimens used to eradicate *H. pylori*, which include broad-spectrum antibiotics and proton pump inhibitors (PPIs). A study showed that a triple-drug combination therapy containing amoxicillin, clarithromycin, and a PPI for 7 days could be a suitable first-line treatment for NHPH as well as *H. pylori* infection.⁸

NHPH species represent an under-recognized group of gastric pathogens with significant clinical implications, yet current studies on NHPH face several limitations. Most of the studies conducted are single-center and have small sample sizes. Also, a significant number of studies had



*Corresponding Author: Amir Mohammad Salehi, Email: amirchsalehi19171917@gmail.com



© 2026 The Author(s). This work is published by Middle East Journal of Digestive Diseases as an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

a retrospective design, which are more prone to biases.⁸ Many aspects of epidemiology, pathogenesis, diagnosis, and proper treatments remain unclear. Therefore, we suggest further comprehensive research to better define the role of NHPH in gastric diseases, improve diagnostic accuracy, and develop an optimal treatment strategy.

Authors' Contribution

Conceptualization: Amir Mohammad Salehi, Maryam Hasanzarrini, Hossain Salehi

Writing—original draft: Romina Rezaei, Alireza Sadeghi

Writing—review & editing: Amir Mohammad Salehi, Maryam Hasanzarrini

Competing Interests

The authors declare no conflict of interest related to this work.

Ethical Approval

There is nothing to be declared.

Funding

Hamadan University of Medical Sciences, Hamadan, Iran, supported this study.

References

1. Taillieu E, De Witte C, De Schepper H, Van Moerkercke W, Rutten S, Michiels S, et al. Clinical significance and impact of gastric non-*Helicobacter pylori* *Helicobacter* species in gastric disease. *Aliment Pharmacol Ther* 2023;57(12):1432-44. doi:10.1111/apt.17488
2. Øverby A, Yamagata Murayama S, Matsui H, Nakamura M. In the aftermath of *Helicobacter pylori*: other helicobacters rising up to become the next gastric epidemic? *Digestion* 2016;93(4):260-5. doi:10.1159/000445399
3. Haesebrouck F, Pasmans F, Flahou B, Chiers K, Baele M, Meyns T, et al. Gastric helicobacters in domestic animals and nonhuman primates and their significance for human health. *Clin Microbiol Rev* 2009;22(2):202-23. doi:10.1128/cmr.00041-08
4. Nakamura M, Øverby A, Michimae H, Matsui H, Takahashi S, Mabe K, et al. PCR analysis and specific immunohistochemistry revealing a high prevalence of non-*Helicobacter pylori* helicobacters in *Helicobacter pylori*-negative gastric disease patients in Japan: high susceptibility to an Hp eradication regimen. *Helicobacter* 2020;25(5):e12700. doi:10.1111/hel.12700
5. Cortez Nunes F, Mateus TL, Aguiéiras C, Louro R, Peixe B, Calhindo M, et al. Prevalence and diagnostic comparison of *Helicobacter pylori* and non-*Helicobacter pylori* *Helicobacter* infections in patients undergoing upper gastrointestinal endoscopy with gastric biopsy in Algarve, Portugal. *Microorganisms* 2025;13(7):1684. doi:10.3390/microorganisms13071684
6. Takeda T, Asaoka D, Hirai S, Nakamura M, Nagahara A. Endoscopic features with linked-color imaging of non-*Helicobacter pylori* *Helicobacter*-associated gastritis: a case study. *VideoGIE* 2024;9(3):123-7. doi:10.1016/j.vgie.2023.10.009
7. Sugimoto M, Rimbara E, Murata M, Yamaoka Y. Roles of gastric non-*Helicobacter pylori* *Helicobacter* species in gastric disease development: a review. *Helicobacter* 2025;30(3):e70051. doi:10.1111/hel.70051
8. Tsukadaira T, Hayashi S, Ota H, Kobayashi N, Agawa H, Kodaira H, et al. Triple-drug combination therapy versus six-month proton pump inhibitor monotherapy in non-*Helicobacter pylori* *Helicobacter* eradication, and hyperacid environment preference of *Helicobacter suis*: a clinical study. *BMC Gastroenterol* 2024;24(1):157. doi:10.1186/s12876-024-03252-5